

# A Review On Dump Slope Stabilization By Revegetation With Reference To Indigenous Plant

Vibhash Ranjan, Arjun Sarsawat, Phalguni Sen, Dheeraj Kumar

**Abstract:** The waste, sub-grade ROM and Fines dumps are characterized by high rock fragment contents, low moisture retention capacity, higher bulk density, low nutrients, lower pH and elevated metal concentrations. Use of suitable revegetation programme, that require the selection of right type of plants to be used vis-à-vis the site condition and characteristics, can enhance the long term stability, both mechanical and ecological, of dumps (waste, sub-grade ROM and fines) through providing vegetative cover to control soil erosion & gully formation, consolidation of dump top & side surfaces, binding the loose particles through intricate root system, etc. This paper overviews the roles of Revegetation programme that can be used where vegetation is considered to be the long term answer to slope protection and erosion control. Various revegetation processes like Coir matting, Lemmon Grass, Vetiver Grass and Indigenous plant etc. may find a wide range of usage in Iron ore mines for dump stabilization. Miyawaki method of plantation is more effective than the conventional method of plantation. This will lead to co-existence of plants and as a matter of fact each plant draws from the other vital nutrients and they grow to become strong and healthy. The Miyawaki Plantation technique aim at "Survival at fittest" the area undergone such plantation have an ecology of their own. The Indigenous Plant has good binding capacity and helps to control soil erosion as well as improve the dump stability. Indigenous plant species like as Shorea robusta (Sal), CROTON ROXBURGHII, Dal-bergia Sisoo (SHISHAM), KARANJ, Azadirachta (NEEM) etc. The growth of indigenous plants is depending upon the soil quality as well as the organic matter of the dump material. The leaf debris plays important role for improving the organic matter of dump material for successfully implementation of revegetation programme.

**Keywords:** Revegetation; Coir Mat; Vetiver Grass; Lemon Grass; Indigenous Plant; Selection of Plant species.

## 1. Introduction

Opencast mining operations involve huge quantities of overburden and sub grade ROM removal, dumping and backfilling of excavated area. Substantial increase in rate of accumulation of waste/sub grade ROM materials in recent years has resulted in greater height of dumps to minimize ground cover area. Consequently it has given rise to the danger of dump failures, gully erosion and various associated environmental problems (Campbell, 1992). Extraction of valuable minerals has been producing wastes in the form of waste rocks whose management is again of prime importance. The issues relating to the stability of these dumps/stacks is catching attention worldwide for quite some time now, which is quite important for the safe working in and around these monstrous structures as well as restricted availability of land. Even government bodies like SPCB, MoEF, and DMG etc. have put raised their concern by way of specified rules and regulations making it mandatory for all mines. Some photographs of critical dump failure in Iron ore mines, (Fig.1-3).



**Fig. 1:** Slope failure in Sub-grade dump



**Fig. 2:** Slope failure in Fines dump

- Vibhash Ranjan, Arjun Sarsawat, Phalguni Sen, Dheeraj Kumar,
- Thakurani Iron Ore Mines of M/s Sarda Mines (P) Limited, Soyabali, Barbil, Keonjhar, (Odisha), India
- Department of Mining Engineering, Indian School of mines, Dhanbad 826004, India, Tel.: +9106767247681; e-mails: vibhash.ranjan@sardamines.net/yes\_raj2001@yahoo.com



**Fig. 3:** Slope failure in Waste dump


The review aims to provide a summation of the issues and information relating to revegetation in waste, sub-grade ROM and fines dump drawing on discussions with regulators and researchers, visits to mine sites throughout the region and a review of the published literature, consultants and government reports. In doing so, the review aims to know the revegetation technique implemented in the different iron ore mines for slope stabilization with respect to their effectiveness. Revegetation is one of the widely used technique for controlling erosion and stabilization of dump slope (Akers and Muter, 1974), and thereby maintaining ecological equilibrium in the area (Jorgensen, 1994). With respect to the Role of vegetation growth upon dump slope can be described as hydrogeological and mechanical actions (Cherubini and Geiasi, 1997) with respect to the hydrogeological action, roots of vegetation play an important role to enhance dump stability by controlling interception of rain water and evapotranspiration and the resulting pore pressure reduction (Hussain, 1995). Whereas mechanical action in turn, reinforced the dump material by roots and enhanced the shear strength of dump material. This action is closely related to root density, depth and strength (Greenway, 1987; Jha, 1989; Suyama, 1992; Hall et al., 1994). The Indigenous Plant like as Sal, Kendu, Amla, Jamun, Shisham Croton roxburghii, Karanj, Neem has good binding capacity and helps to control soil erosion as well as improve the dump stability. Four species (Karanj, *Pongamia* sp.; Shisham, *Dalbergia* sp.; Shirish, *Albizia* sp.; and, Neem, *Azadirachta* sp.) have emerged as having the highest frequency of reporting in addition to an array of shared and relevant agro- and socioeconomically relevant properties along with notably similar environmental outcomes regarding their impact toward above- and belowground ecological factors. These characteristics include that three (members of the Fabaceae) are promoters of nitrogen fixing interactions, while all four are relatively fast growers, well adapted to environmental conditions within more arid zones (i.e., intense heat, sunlight), and have root architectural adaptations for drought tolerance (Koul et al. 1990; Lowry et al. 1994; Tewari 1994). The effectiveness of revegetation can be measured through Numerical modelling in terms of factor of safety before and after implementation of revegetation programme

## 2. Revegetation Process and Establishment

Revegetation process is one of the important methods which can make the ecologically challenging task of restoring the previously existing ecosystem. Revegetation is normally implemented during the dormant season, usually monsoon or winter. Constraints on planting times and/or availability of suitable plant materials may limit the usefulness of certain methods. Select plant materials that are adaptive to the site conditions. Local species are readily available and well suited to the climate, soil condition, and available moisture; therefore they make good candidates. The Indigenous Plant species like as Sal, Kendu, Amla, Jamun, Shisham Croton roxburghii, Karanj, Neem have good binding capacity and helps to control soil erosion as well as improve the dump stability. The evolution of mine revegetation practices has seen the agronomic engineering approach of establishing 'green' landscapes with introduced grass and pasture species (Dahl & Mulligan 1996) replaced by the re-establishment of native flora to create a self-sustaining ecosystem. An early attempts to revegetate mined areas at Weipa and Rum Jungle relied on introduced grass and tree species and nitrogen-fixing shrubs (Dahl & Mulligan 1996, Richards et al 1996).

### Coir Matting

Coir mats and netting are dense, biodegradable fabrics made of coconut husk fiber (coir). They are used to trap sediment, protect stream banks and wetland shores from erosion, and to provide a stable substrate for plant establishment. Coir mat is a kind of natural Erosion Control Blanket (Figs.1–2).



**Fig.1:** Coir Mat



**Fig. 2:** Installation of Coir Mat

## CASE REFERENCES

Erosion Control blankets like Coir mats have been used for slope protection at many places specially in coal mines. Coir mats have also been used to protect high overburden dumps of fly ash and it has been possible to establish vegetation over the dump. An interesting case study where the effectiveness of Gabions and Coir mat has been successfully implemented is the stabilization of overburden dumps at coal mines of Singrauli. A huge dump of 120 m high could be stabilized by combination of Gabion walls and Coir mats on slopes. Of course an in-depth study of strata along with stability analysis combined with quality work at site, all in collection has contributed to the success of the project.

## Disadvantage


It is not fully successfully implemented in the iron ore mines of Odisha state due to the following reasons.

- Poor binding properties with the different size of dump materials
- It requires huge quantity of water during the seeding process till plantation
- Not a suitable option for large and steep slope of dump area.
- There is no use of end product.
- Poor stabilizing qualities rather than Indigenous plant.
- Poor stabilization of failure surface during heavy rain & wind due to its poor binding properties.

**Few observations:-** (Photographs of typical failure of coir matting in the iron ore mines of Joda circle in Odisha (Fig. 1 to 2).




(Fig.1)



(Fig.2)

## Vetiver Grass

The indigenous knowledge was taken along when Indians migrated overseas and usage resumed in new localities around the globe. Thus one reads of vetiver usage for slope protection and reinforcement of embankments and cuttings outside of farmlands since the early 1900's in the West Indies, South Africa (NRC, 1993), Brazil (Grimshaw, 1994) and Fiji (Truong and Gawander, 1996), etc. In 1931, it was on record that vetiver was grown at Serdang (near Kuala Lumpur), Malaysia, where "it is used for holding up steep banks. It is well known to be good for this purpose" (World Bank, 1995). These applications of vetiver, however, were based on past experiences that showed them to be effective but without any quantitative engineering data to back them up. To lend technical credence to its introduction, researches have been conducted on the roles of vegetation in relation to slope stability, in particular on tree roots which the main factor are contributing to slope stabilization (Greenway, 1987; Coppin and Richards, 1990; Gray, 1994;). Vetiver Grass Technology (VGT-Fig 1 to 2) was first



(Fig.1)




Fig. 2 (Root system)

developed for soil and water conservation in farmlands. While this application still plays a vital role in agricultural lands, vetiver grass's unique morphological, physiological, and ecological characteristics have a key role in the area of environmental protection. Unique morphological characteristics include a massive, finely structured, deep root system capable of reaching 2 to 3 meter. The Vetiver Grass Technology (VGT), in its most common form, is simply the establishment of an arrow (less than 1

meter wide) live stiff vetiver grass barrier, in the form of a hedge, across the slope of the land. When applied correctly the technology is effective on slopes from less than 1 to over 100%. In addition its massive root system will increase the shear strength of soil (thus providing improved stability of soils on steep slopes). Vetiver grass is a clump grass, with erect and stiff stems that grow to as much as 2m high.

#### Disadvantage:-

- Seed easily spread in the wind- may become a weed, infesting an area
- Unpalatable to livestock (unsuitable fodder)
- Not suitable for human consumption
- The root of vetiver grass is growing vertically not in a radial direction.
- Oil extraction unlikely option for poor farmers as an end product
- Lower stabilizing qualities rather than Indigenous trees.
- Poor stabilization of failure surface due to its vertical roots.

#### Lemon Grass

An aromatic grass species has indicated that lemon grass (*Cymbopogon citrates*) can be considered as one of the options for the slope stabilization and control of soil erosion. Lemon grass (*Cymbopogon citrates*, is one of the species that one can use for commercial production) is a tropical grass type originally from India. It derives its name from its strong fresh lemon scent. It is a tufted perennial grass with numerous stiff leafy stems arising from short rhizomatous roots. The leaves can grow up to a height of 2m. It is a perennial crop and has an economic lifespan for about 5 years. The plant grows on a wide range of soils. The best yields will be obtained from well drained sandy to loam soils, with a pH value ranging from 4.3 to 8.4 and with access to water throughout the year, it can give 3 - 4 harvests. Often the grass is grown on poor soils or erosion sensitive areas. Without water it will, through its deep rooting system, survive until the next raining season. The root of lemon grass is uniaxial direction rather than vertical. It has been rarely implemented in the dump slope area in iron ore mines.

Few photographs of implementation of Lemon Grass in ground area (Herbal Park) (Fig.1-2)



(Fig.1)



(Fig.2)

#### Disadvantage:-

- Seed easily spread in the wind- may become a weed, infesting an area
- It is most suitable in soil contained material.
- It requires huge quantity of water during the plantation
- Not a suitable option for small holdings of hilly areas
- Lower stabilizing qualities rather than Indigenous plant.
- Poor stabilization of failure surface due to its binding properties.
- It is most prone to fire during dry season.

Further, Use of Grass-Legume mixtures to stabilize dump slopes of sponge iron plant has now become a widely used technique. Studies have shown that *Pennisetum Pedicellatum* along with a forage legume *Styosanthes humilis* can successfully restore a degraded site and stabilization the dump slope of sponge plant (Maiti, S. k, Saxena, N.C. 1998; Araujo, I.C.S., Costa, M.C.G, 2013).

The roots of smaller plants (such as grasses and shrubs) do not go deep and therefore, the overall effect of grass on dump slope is negligible. However, they stabilize the slope by binding the upper layer of dump slope. It is observed that the small plants do not significantly improve the factor of safety (FOS) of slope. However, it could be useful for early stabilization. The grasses quickly bind the upper surface, whereas shrubs too immensely strengthen the stability of the dump in the initial stage. Further it was concluded that the Trees like (Sisam-group of indigenous plant) for long term stability as the roots of trees are stronger and penetrate deeper. Their roots bind the dump material and form a composite material having high shear strength. The Sisam tree of even moderate size binds the upper layers of mine dump and improve the factor of safety substantially. It was also observed that the stability of dump slope improves with respect to time i.e. size of tree (Rai Rajesh, Srivastava B.K, 2010 & 2014).

### 3. Revegetation as a stabilizing agent on dump slopes

There is a generalized agreement that "forest plays an important role in the protection of soil, and deforestation can bring about not only erosion, but also collective soil movements"(Gray, 1973). Sternberg (1975) mentions studies by bailey, who by measuring the maximum angle of

the repose of slopes with and without vegetation, found that a given soil derived from the same rock pattern, the maximum angle of repose is about 60° when vegetation exists and about 36°, in the case of a lack of vegetation.

Vegetation has an important role in protecting the soil surface from erosion and allowing accumulation of fine particles (Tordoff et al., 2000; Conesa et al., 2007b). They can reverse degradation process by stabilizing soils through development of extensive root systems. Once they are established, plants increase soil organic matter, lower soil bulk density, and moderate soil pH and bring mineral nutrients to the surface and accumulate them in available form. Their root systems allow them to act as scavengers of nutrients not readily available. The plants accumulate these nutrients and redeposit them on the soil surface in organic matter from which nutrients are much more readily available by microbial breakdown (Li, 2006; Conesa et al., 2007a).

The revegetation of eroded ecosystems must be carried out with plants selected on the basis of their ability to survive and regenerate or reproduce under severe conditions provided both by the nature of the dump material, the exposed situation on the dump surface and on their ability to stabilize the soil structure (Madejon et al., 2006). Normal practice for revegetation is to choose drought-resistant, fast growing crops or fodder which can grow in nutrient deficient soils. Selected plants should be easy to establish, grow quickly, and have dense canopies and root systems. In certain areas, the main factor in preventing vegetation is acidity. Plants must be tolerant of metal contaminants for such sites (Caravaca et al., 2002; Mendez and Maier, 2008b). Role of exotic or native species in reclamation needs careful consideration as newly introduced exotic species may become pests in other situations. Therefore, candidate species for vegetation should be screened carefully to avoid becoming problematic weeds in relation to local to regional floristic. For artificial introduction, selection of species that are well adapted to the local environment should be emphasized. *Indigenous species are preferable to exotics* because they are most likely to fit into fully functional ecosystem and are climatically adapted (Chaney et al., 2007). Grasses are considered as a nurse crop for an early vegetation purpose. Grasses have both positive and negative effects on mine lands. They are frequently needed to stabilize soils but they may compete with woody regeneration. Grasses, particularly C4 ones, can offer superior tolerance to drought, low soil nutrients and other climatic stresses. Roots of grasses are fibrous that can slow erosion and their soil forming tendencies eventually produce a layer of organic soil, stabilize soil, conserve soil moisture and may compete with weedy species. The initial cover must allow the development of diverse self-sustaining plant communities (Shu et al., 2002; Singh et al., 2002; Hao et al., 2004). Trees can potentially improve soils through numerous processes, including - maintenance or increase of soil organic matter, biological nitrogen fixation, uptake of nutrients from below and reach of roots of under storey herbaceous vegetation, increase water infiltration and storage, reduce loss of nutrients by erosion and leaching, improve soil physical properties, reduce soil acidity and improve physical properties, reduce soil acidity and improve soil biological activity. Also, new self-sustaining top soils are created by trees. Plant litter and root exudates provide nutrient-cycling to soil (Pulford and Watson, 2003; Coates,

2005; Padmavathamma and Li, 2007; Mertens et al., 2007). On mine spoils, nitrogen is a major limiting nutrient and regular addition of fertilizer nitrogen may be required to maintain healthy growth and persistence of vegetation (Yang et al., 2003; Sing et al., 2004). An alternative approach might be to introduce legumes and other nitrogen-fixing species. Nitrogen fixing species have a dramatic effect on soil fertility through production of readily decomposable nutrient rich litter and turnover of fine roots and nodules. Mineralization of N-rich litter from these species allow substantial transfer to companion species and subsequent cycling, thus enabling the development of a self-sustaining ecosystem (Zhang et al., 2001). Singh et al., (2002) reported that native leguminous species show greater improvement in soil fertility parameters in comparison to native non-leguminous species. Also, native legumes are more efficient in bringing out differences in soil properties than exotic legumes in the short term.

**4. Selection of Plant Species-** *a vital component for eco restoration success.* The choice of plant species depends on the characteristics of spoil, climatic condition and ultimate land use. Indigenous plant are like as Sal, Kendu, Amla, Jamun, Shisham Croton roxburghii, Karanj, Neem etc. are more tolerant to adverse spoil pH and moisture stress than grass, legumes and are therefore, easier to establish. This indigenous plant are used where a wind break is desired or where visual shield is required they are suitable to steep or rocky terrain and coarse waste. The successful vegetation establishment of waste dumps also depends on micro and macro-climate, disease and insect resistance, competition, growth pattern and propagation ability of the species. The availability of moisture in the spoil, especially during warm dry summer, is important. If the water level is not high enough, irrigation may be necessary to provide moisture. As the vegetation matures the need for the irrigation diminishes since Indigenous plant gets adapted to dry summer condition. In spite of the relatively modest peer-review literature pool available on the topic of post-mined land revegetation in India, there is still a remarkably extensive reporting of plant species (n=245 sp.) including vegetative growth forms and diverse bioindicators of ecological form and function. With this rationale, four species (Karanj, *Pongamia* sp.; Shisham, *Dalbergia* sp.; Shirish, *Albizia* sp.; and, Neem, *Azadirachta* sp.) have emerged as having the highest frequency of reporting in addition to an array of shared and relevant agro- and socioeconomically relevant properties along with notably similar environmental outcomes regarding their impact toward above- and belowground ecological factors. These characteristics include that three (members of the Fabaceae) are promoters of nitrogen fixing interactions, while all four are relatively fast growers, well adapted to environmental conditions within more arid zones (i.e., intense heat, sunlight), and have root architectural adaptations for drought tolerance (Koul et al. 1990; Lowry et al. 1994; Tewari 1994). The growth of indigenous plants is depending upon the soil quality as well as the organic matter of the dump material. The leaf debris plays important role for improving the organic matter of dump material for successfully implementation of revegetation.

## 5. Factors to consider in selecting species for revegetation

According to Morgan (2005), species to be used for revegetation should be selected based on their following parameters; properties of rapid growth; toughness in respect of diseases and pests; ability to compete with less desirable species; adaptability to the local soil; and adaptability to the local climatic conditions. When the species to be used in the revegetation programmes are trees, they should possess the following qualities, according to Young (1997), for it to be qualified for use in soil fertility improvement/restoration: Primarily have the capacity to grow on poor soils; High rate of nitrogen fixation; High and balanced nutrient content in foliage litter; Litter low in lignin and polyphenols; Either rapid litter decay where nutrient release is desired or moderate rate of litter decay where soil cover is required; Absence of toxic substances in litter or root residues; Well-developed tap root system; High rate of leafy biomass production; and dense network of fine roots with a capacity for mycorrhizal association. Species selection will depend on local soil and climatic conditions but should aim to provide a uniform rather than clumpy pattern of vegetation to avoid concentrations of runoffs and localized erosion. The ideal condition is difficult to achieve, however, where the soils have low water retention capacity or are toxic, or where cold, drought or exposure inhibits plant growth. Choosing vegetation which has beneficial effect is important (Morgan, 2005). For instance, in order to reduce the cost of reclamation of mine spoil mounds from tin mining on the Jos Plateau, Nigeria, the Mine Lands Reclamation Unit chose *Eucalyptus* sp. which could be cropped as a source of fuel and pole timber but these have resulted in significant declines in the base status and pH of the soil (Alexander, 1990) to the detriment of other plant growth (Morgan, 2005). The revegetation of eroded ecosystems must be carried out with plants selected on the basis of their ability to survive and regenerate or reproduce under severe conditions provided both by the nature of the dump material, the exposed situation on the dump surface and on their ability to stabilize the soil structure (Madejon et al., 2006). Normal practice for revegetation is to choose drought-resistant, fast growing crops or fodder which can grow in nutrient deficient soils. Selected plants should be easy to establish, grow quickly, and have dense canopies and root systems. In certain areas, the main factor in preventing vegetation is acidity (Sheoran et al., 2010). Plants must be tolerant of metal contaminants for such sites (Caravaca et al., 2002; Mendez and Maier, 2008b).

## 6. Conclusion

For centuries, wood and indigenous living plants were the only material for hill and slope stabilization works. With advancing technology, this age old concept has been further refined and some innovative techniques are being adopted. In the wake of alarming situation of Global warming, it is highly encouraged to adopt ecofriendly solutions and promote the growth of nature. Revegetation is the most appropriate for environmentally and aesthetically sensitive areas. Various Ecological Restoration approach like Coir matting, Lemmon Grass, Vetiver Grass and Indigenous plant etc. may find a wide range of usage in Iron ore mines for dump slope stabilization. The roots of smaller plants (such as grasses and shrubs) do not go deep and

therefore, the overall effect of grass on dump slope is negligible. However, they stabilize the slope by binding the upper layer of dump slope. It is observed that the small plants do not significantly improve the factor of safety (FOS) of slope. However, it could be useful for early stabilization. The grasses quickly bind the upper surface, whereas shrubs too immensely strengthen the stability of the dump in the initial stage. Further it was concluded that the Trees like (Sisam-group of indigenous plant) for long term stability as the roots of trees are stronger and penetrate deeper. Their roots bind the dump material and form a composite material having high shear strength. The Sisam tree of even moderate size binds the upper layers of mine dump and improve the factor of safety substantially. A further study is required in respect of dump slope stabilization with reference to Indigenous Plant. Indigenous plant is a native typical dry deciduous forest is dominated by tree species *Shorea robusta* (Sal), *CROTON ROXBURGHII*, *Dal-bergia sisoo* (SHISHAM, KARANJ, Azadirachta ( NEEM )having good binding capacity and helps to control soil erosion as well as improve the dump stability. The growth of indigenous plants is depending upon the soil quality as well as the organic matter of the dump material. It is pertinent to mention that different parts of a plant perform specific functions. Roots increase hydraulic conductivity of soil and reinforce them by performing functions of anchorage, absorption, conduction and storage. Stems help in interception. Leaves aid storage and add to the beauty. In fact vegetation helps dissipate the kinetic energy of rain drops, attenuates the effects of wind and helps reduce the velocity of overland flow. The most critical aspect of revegetation intervention is the choice of plants and vegetation. But the problem needs to be properly analyzed and the technically best possible solution needs to be adapted without any compromise to safety of the structure and surroundings. In order to get complete advantage of the environmental friendly solution, it is best to select the indigenous plants with the help of a horticulturist after studying the area, its climatic conditions, humidity range, soil type, annual rainfall and other relevant criteria

## 7. Bibliography

- [1] Akers, J.D., Muter, (1974). Gob pile stabilization and reclamation. Proceeding of the Forth Mineral Waste Utilization Symposium, Chicago, Illinois, pp. 229-239
- [2] Alexander MJ (1990). Reclamation after tin mining on the Jos plateau, Nigeria. *Geographical J.* 156:44-50
- [3] Araujo, I.C.S., Costa, M.C.G. "Biomass and nutrient accumulation pattern of leguminous tree seedlings grown on mine tailing amended with organic waste". *Ecol Eng.*60, pp. 254-260, (2013).
- [4] Campbell, D.B. (1992). Resloping of waste rock dumps. *Int.Mine Waste Manag.News* 2 (2), 7-10
- Cherubini, C., Geiasi, C., (1997). The influence of vegetation on slope stability. In: Marinos, P.G., Koukias, G.C., Tsiambaos, G.C., Stournaras, G.C. (Eds.), *Engineering Geology and Environment*, vol. I. Balkema, Rotterdam, pp. 67-71.

- [5] Chaulya, S. K., Singh, R. S. and Dhar, B. B. (1999). Growth performance of selected grass species on coal-mine overburden dump slopes in India, *Trans. Instn. Min. Metall. (Sect. A: Min. Industry)*, 108, A136–139
- [6] Caravaca, F., Hernandez, M.T., Garcia, C., and Roldan, A. (2002). Improvement of rhizosphere aggregates stability of afforested semi- arid, plant species subjected to mycorrhizal inoculation and compost addition. *Geoderma* 108,133-144
- [7] Chaney, R.L., Angle, J.S., Broadhurst, C.L., Peters, C.A., Tappero, R.V., and Donald, L.S. (2007). Improved understanding of hyperaccumulation yields commercial phytoextraction and phytomining technologies. *Journal Environmental Quality* 36, 1429-14423.
- [8] Coppin NJ, and Richards IG. (1990). Use of vegetation in civil engineering construction industry and research information association (CIRIA), Butterworths, London.
- [9] Conesa, H.M., Schulin, R., and Nowack, B. 2007a. A laboratory study on revegetation and metal uptake in native plant species from neutral mine tailings. *Water, Air, and Soil Pollution*, 183 (1-4), 201-212
- [10] Conesa, H.M., Garcia, G., Faz, A., and Arnaldos, R. 2007b. Dynamics of metal tolerant plant communities' development in mine tailings from the Cartagena-La Union Mining District (SE Spain) and their interest for further revegetation purposes. *Chemosphere* 68, 1180-1185.
- [11] Dahl NW & Mulligan DR (1996). Environmental management at Weipa: Bauxite mining in the tropical north. In *Environmental management in the Australian minerals and energy industry: Principles and practices*, ed DR Mulligan, UNSW Press, Sydney, 403–420
- [12] Gray, D.H. (1973) Effects of forest clear cutting on the stability of natural slopes: results of field studies. National Science Foundation, University of Michigan, Washington
- [13] Gray DH. (1994). Influence of vegetation on stability of slopes. In: *Vegetation and Slopes*. Institution of Civil Engineers, London. 2-25
- [14] Greenway D R (1987), "Vegetation and Slope Stability, Slope Stability", in M G Anderson and K S Richards (Eds.), *Slope Stability*, John Wiley and Sons Ltd. New York
- [15] Grimshaw RG. (1994). Vetiver grass—its use for slope and structure stabilization under tropical and semitropical conditions. In: *Vegetation and Slopes*. Institution of Civil Engineers, London. 26-35
- [16] Hall B E, Giles E L and Rauch H P (1994), "Experiences with the Use of Trees in Slope Stabilization", in *Proceedings of XIII ICSMFE*, pp. 1231-1235, New Delhi
- [17] Hussain, A., (1995). Fill compaction-erosion study in reclaimed areas. *Indian Mining Eng.*J.34 (6), 19-21
- [18] Hao, X. Z., Zhou, D. M., Wang, Y. J., and Chen, H.M. (2004). Study of rye grass in copper mine tailing treated with peat and chemical fertilizer. *Acta Pedol Sin* 41(4), 645-648
- [19] Jha AK (1989), "A note on the Root Development in Dry Tropical Naturally Revegetated Coal Mine Spoil", *Vegetation*, Vol. 85, pp. 67-70.
- [20] Jorgensen, S.E., (1994). Models as instruments for combination of ecological theory and environmental practice. *Ecol.Model.* 75/76, 5-20.
- [21] Koul O, Isman MB, and Ketkar CM (1990) Properties and uses of neem, *Azadirachta indica*. *Canadian Journal of Botany* 68, 1-11
- [22] Li, M.S. (2006). Ecological restoration of mineland with particular reference to the metalliferous mine wasteland in China: a review of research and practice. *Soil Total Environment* 357, 38-53
- [23] Lowry JB, Prinsen H, and Burrows DM (1994) *Albizia lebbeck*: a promising forage tree for semiarid regions. In „Forage Tree Legumes in Tropical Agriculture“. (Ed Gutteridge RC and Shelton HM) pp 75-83. (CAB International: Wallingford, Oxon, UK).
- [24] Maiti, S.k.Saxena, N.C., "Biological reclamation of coal mine spoils without topsoil: an amendment study with domestic raw sewage and grass legumes mixtures". *Int. J. Surf. Min. Reclamat Environ.*12, pp.87-90, (1998)
- [25] Morgan RPC (2005). *Soil erosion and Conservation* (3rd edition). Blackwell Publishing Company Ltd, U.K. 59; 169-70; 190-91
- [26] Madejon E, de Mora AP, Felipe E, Burgos P, Cabrera F (2006). Soil amendments reduce trace element solubility in a contaminated soil and allow re-growth of natural vegetation. *Environ. Pollute.* 139: 40-52
- [27] Mendez, M.O., and Maier, R.M. 2008b. Phytostabilization of mine tailings in arid and semiarid environments-An emerging remediation technology. *Environmental Health Perspectives* 116 (3), 278-283.
- [28] National Research Council. (1993). *Vetiver Grass: A Thin Green Line against Erosion*. National Academy Press, Washington DC, USA

- [29] Pulford, I. D., and Watson, C. (2003). Phytoremediation of heavy metal-contaminated land trees-a review. *Environmental International* 29, 529-540.
- [30] Padmavathiamma, P.K., and Li, L.Y. (2007). Phytoremediation technology: Hyperaccumulation metals in plants. *Water Air Soil Pollution* 184(1-4), pp.105-126
- [31] Rai Rajesh and Shrivastava B.K (2011) Biological stabilization of mine dumps: Shear strength and numerical simulation approach with reference to Sisam Tree, *Environ Earth Sci* (2011) 63:177–188.
- [32] Rai Rajesh and Shrivastava B.K (2014) Numerical simulation of vegetated mine dump slope with reference to small plants, *International Journal of Mining Science and Technology* 24 (2014) 111–115.
- [33] Richards RJ, Applegate RJ & Richie AIM (1996). The Rum Jungle rehabilitation project. In *Environmental management in the Australian minerals and energy industries*. ed DR Mulligan, UNSW Press & AMEEF, Sydney, 464–553.
- [34] Sternberg, H.O. (1975) Effects of deforestation on slopes. *Journal of the Geotechnical Engineering Division, ASCE* 101: pp.147-165.
- [35] Singh, A.N., Raghubanshi, A.S., and Singh, J.S. (2002). Plantations as a tool for mine spoil restoration. *Current Science* 82(12), 1436–1441
- [36] Shu, W.S., Xia, H.P., Zhang, Z.Q., and Wong, M.H. (2002). Use of vetiver and other three grasses for revegetation of Pb/Zn mine tailings: field experiment. *International Journal of Phytoremediation*, 4(1): 47-57.
- [37] Singh AN, Raghubanshi AS, Singh JS (2004) Impact of native tree plantations on mine spoils in a dry tropical environment. *Forest Ecology and Management* 187:49–60
- [38] Sheoran V, Sheoran A S, Poonia P (2010). Soil Reclamation of Abandoned Mine Land by Revegetation: A Review, *Intl J. Soil, Sediment and Water*. 3: 1-21. Available at: <http://scholarworks.umass.edu/intljssw/vol3/iss2/13>
- [39] Suyama M (1992), "Assessment of Biotechnical Slope Stability Effect for Urban Forest in Japan", in *Proceedings of 6th International Symposium on Landslides*, pp. 831-836, A A Balkema, Christchurch.
- [40] Tewari DN (1994) (Ed) „A monograph on Dalbergia sissoo Roxb.“ (Ed Tewari DN) (International Book Distributors: Dehradun, India).
- [41] Truong P, and Gawander JS. (1996). Back from the future: do's and don'ts after 50 years of vetiver utilization in Fiji. *Proceedings of the First International Conference on Vetiver*. Office of the Royal Development Projects Board, Bangkok. 12-17
- [42] Tordoff, G.M., Baker, A.J.M., Willis, A.J., 2000. Current approaches to the revegetation and reclamation of metalliferous mine wastes. *Chemosphere* 41, 219-228.
- [43] Young A (1997). *Agroforestry for Soil Management, Effects of trees on soils*. CAB International. 23
- [44] Yang, B., Shu, W. S., Ye, Z. H., Lan, C. Y., and Wong, M. H. (2003). Growth and metal accumulation in vetiver and two *Sesbania* species on lead/zinc mine tailings. *Chemosphere* 52(9), 1593-1600.
- [45] Zhang ZQ, Shu WS, Lan CY, Wong MH (2001). Soil seed bank as an input of seed sources in vegetation of lead/ zinc mine tailings. *Restoration Ecol.* 9(4):1-8.